Seizure Emergency Care Plan and Medication Orders for School and Childcare Settings	
PARENT/GUARDIAN complete and sign the	
	Birth date:
	Phone: Place
<u> </u>	Phone: child's
	Grade: photo here
Triggers:  tiredness flashing lights illness hunger temperature Other: Seizure Aura (if any):	
Seizure history: Convulsive Focal Absence Date of last known seizure  Describe:	
Antiseizure Medication Taken at Home Common	side effects
Other Seizure Treatments/Special Diet Therapy:	
I give permission for school personnel to share this information, follow this plan, administer medication and care for my child and, if necessary, contact our health care provider. I assume full responsibility for providing the school with prescribed medication and devices. I approve this Seizure Emergency Care Plan for my child.	
PARENT SIGNATURE DATE	SCHOOL NURSE SIGNATURE DATE   IEP
HEALTH CARE PROVIDER to complete all items.  IF YOU SEE THIS:	, SIGN and DATE completed form.  DO THIS:
Convulsive Generalized Tonic Clonic:	Time the seizure
You will see loss of consciousness. Stiffening of the bod' Rhythmic jerking movements. Convulsive seizures may last 1-5 minutes. The child may have a warning (aura) before the seizure. Sleepiness and confusion may occur after the seizure.	<ol> <li>Keep calm. Provide reassurance.</li> <li>Protect head, keep airway clear, turn on side if possible.</li> <li>Do not place anything in mouth.</li> </ol>
☐ Focal: These seizures may begin with an aura. They may be partly alert or unconscious. You may see lip smacking, chewing, eye blinking, or picking at clothes. These seizur usually last 1-2 minutes.  ☐ Absence: You will see quick changes in alertness. May see eye flutter or small twitching. Usually last less than 10 seconds.	Time the seizure     Gently guide child away from danger.     Stay with student and reassure them until recovered from seizure.     Do not treat staring that is stopped by a touch or a nudge.     Call parent.     Administer rescue treatments as marked below.
Rescue Treatments	
☐ Child has a VNS. Child/staff may swipe with aura. Staff may swipe at onset of seizure and every 60 seconds until seizure stops.  Give rescue medications below if seizure does not stop withinminutes.	
If seizure lasts longer than minutes administer:	
☐ Diastatmg rectally ☐ Midazolammg in the nose ☐ Clonazepammg in the cheek ☐ Midazolammg in the nose ☐ Clonazepammg in the cheek ☐ Diastatmg in the cheekmg in the cheek	
☐ Multistep seizure rescue plan − Please see attached letter for details.	
If <u>cluster</u> of or more seizures in min administer:	
☐ Diastatmg rectally ☐ Midazolammg in the nose ☐ Clonazepammg in the cheek ☐ Multistan asizura resource plane. Planes are attached latter for details.	
☐ Multistep seizure rescue plan – Please see attached letter for details.	
If emergency medication is administered:  Call 911 immediately or Call 911 if seizure does not stop within 5 minutes Other:	
If no emergency medication is at school and the child is experiencing seizures:	
Call family to bring medications to school or pick up child. Call EMS if seizure lasts more than min	
<b>Accommodations</b> : Always take seizure action plan and emergency medication for school activities, sports and field trips. Close adult supervision when swimming or climbing.	
HEALTH CARE PROVIDED SIGNATURE PRINTS	DROVIDED'S NAME PHONE/EAY DATE